

Sheep Anti-Human Antithrombin (ATIII) Whole IgG from antiserum (10 mg)

Catalogue: SAAT-IG

Lot No: SAMPLE

Expiry date: **/****

Store at -10 to -20°C.

For Research Use Only.

Not for use in diagnostic procedures.

Description of Antithrombin (ATIII)

Antithrombin, also known as Antithrombin III (ATIII), is a member of the SERPIN family of proteinase inhibitors and the primary inhibitor of thrombin in plasma. It is produced in the liver and circulates in plasma at ~200 µg/mL (~3.5 µM). Antithrombin inhibits a broad spectrum of serine proteases including thrombin, activated forms of factor X, factor IX, factor XII, as well as kallikrein, plasmin and urokinase. Enzyme inhibition by antithrombin occurs through proteolytic cleavage at Arg³⁸⁵-Ser³⁸⁶ and subsequent rapid formation of a stable, inactive 1:1 enzyme-antithrombin complex. Heparin has a profound accelerating effect on the inhibitory activity of antithrombin towards some enzymes. For example, the rate of inhibition of thrombin and activated factor X is increased 1000-fold in the presence of optimal concentrations of heparin, whereas heparin has relatively little effect on the inhibition rate of activated factor XI, activated factor XII and kallikrein. Antithrombin is a single chain molecule with a molecular weight of 59 kDa. Interaction with thrombin results in an SDS-stable thrombin-antithrombin complex of 96 kDa¹⁻³.

References and Reviews

1. Damus PS, Rosenberg RD; Antithrombin - Heparin Cofactor; Methods in Enzymology **45**, pp 653-669, 1976.
2. Harpel PC; Blood Proteolytic Enzyme Inhibitors: Their Role in Modulating Blood Coagulation and Fibrinolytic Enzyme Pathways; in Hemostasis and Thrombosis, eds. RW Colman, J Hirsh, VJ Marder and EW Salzman, pp. 738-747, J.B Lippincott Co., Philadelphia PA, USA, 1982.
3. Griffith MJ, Noyes CM, Church FC; Reactive Site Peptide Structural Similarity between Heparin Cofactor II and Antithrombin III; JBC: **260**, pp 2218-2225, 1985.

Product Specifications

Description:

Vial containing 1.000 ml of whole IgG representing approximately 1 ml of antiserum. Total protein is 10 mg.

Format:

Whole IgG, clear liquid.

Host Animal:

Sheep

Immunogen:

Human antithrombin purified from plasma.

Concentration:

IgG concentration is 10 mg/mL, determined by absorbance using an extinction coefficient ($E_{280}^{1\%}$) of 13.4.

Buffer:

10 mM HEPES, pH 7.4, 150 mM NaCl, 50% (v/v) glycerol.

Storage:

Store between -10 and -20 °C. Product will become viscous but will not freeze. Avoid storage in frost-free freezers. Keep vial tightly capped. Allow product to warm to room temperature and gently mix before use.

Specificity:

This antibody is specific for antithrombin as demonstrated by immunoelectrophoresis and ELISA.

Applications:

Suitable as a source of antibodies to Human antithrombin.

Neutralizing activity:

Not determined.

Related Products

GAAT-IG	Goat Anti-ATIII, whole IgG from antiserum
GAAT-AP	Goat Anti-ATIII, affinity-purified IgG
SAAT-AP	Sheep Anti-ATIII, affinity-purified IgG
SAAT-APHRP	Sheep Anti-ATIII, APIgG-peroxidase
ATIII-EIA	Paired antibody set for ELISA of ATIII, 5 x 96 wells
TAT-EIA	Paired antibody ELISA of Thrombin-ATIII complexes
AT-NU10	Paired antibody set for ATIII neutralization in plasma
ATIII-DP	Human plasma deficient in ATIII, immune depleted
ATHC-DP	Human ATIII & HCII def plasma, immune depleted

Visit our site (www.enzymeresearch.co.uk) for further details.

Enzyme Research Laboratories Ltd
Glamorgan House . Charter Court . Enterprise Park .
Swansea . SA7 9DD . UK.
Tel: +44 (0)1792 686070
Email: info@enzymeresearch.co.uk
Website: www.enzymeresearch.co.uk