



Sheep Anti-Human Protein C (PC) Peroxidase Conjugated IgG 0.2 mg

Catalogue: SAPC-HRP
Lot No: SAMPLE
Expiry date: **/****
Store at -10 to -20°C.

For Research Use Only ~ Not for use in diagnostic procedures.

Description of Protein C (PC)

Protein C (PC) is a vitamin K-dependent glycoprotein produced in the liver. The concentration of PC in plasma is ~4 µg/mL (~60 nM). A deficiency of Protein C (quantitative or qualitative) is a risk factor for vascular thrombosis. Protein C is expressed as a two-chain molecule with a molecular weight of 62 kDa. The light chain (21 kDa) of PC consists of two EGF-like domains and an amino-terminal domain containing one hydroxyaspartic acid and 11 γ-carboxyglutamic acid (gla) residues. These residues allow PC to bind to membranes that contain acidic phospholipids in a calcium dependent manner. The heavy chain of PC (41 kDa) consists of the catalytic domain and an activation peptide. Activation of Protein C results from cleavage at residue Arg¹² in the heavy chain by a complex of thrombin and a cell surface cofactor thrombomodulin. The activation of PC is associated with the release of a small activation peptide (2-3 kDa, called Protein C peptide, or PCP) from the N-terminal of the heavy chain. Activated Protein C (APC) is a serine protease with anticoagulant activity. APC, in complex with a phospholipid membrane, calcium and the Protein S cofactor, exhibits anticoagulant activity through the proteolytic inactivation of coagulation cofactors Va and VIIIa. The primary inhibitor of APC activity in plasma is Protein C Inhibitor (PCI, also called Plasminogen Activator Inhibitor-3, PAI-3) and to a lesser extent by α₁-antitrypsin and α₂-macroglobulin. The inhibitory activity of PCI is stimulated approximately 10 fold by heparin¹⁻³.

References and Reviews

1. Broze GJ, Miletich JP; Biochemistry and Physiology of Protein C, Protein S and Thrombomodulin; in Haemostasis and Thrombosis, 3rd Edition, eds. RW Coleman, J Hirsh, VJ Marder and EW Salzman, pp 259-276, J.B. Lippincott Co., Philadelphia PA, USA, 1994.
2. Esmon CT, Esmon NL, Le Bonniec B, Johnson AE; Protein C Activation; Methods in Enzymology 222, pp 359-385, 1993.
3. Heeb MJ, Mosher D, Griffin JH; Activation and Complexation

of Protein C and Cleavage and Decrease of Protein S in Plasma of Patients With Intravascular Coagulation; Blood 73, pp 455-461, 1989.

Product Descriptions

Description:

Vial containing 0.100 mL of whole IgG conjugated to horseradish peroxidase (HRP) through carbohydrate groups. Total protein is 0.2 mg.

Format:

IgG-HRP conjugate as a clear, slightly red-brown liquid.

Host Animal:

Sheep

Immunogen:

Human Protein C purified from plasma.

Concentration:

IgG concentration is 2.00 mg/ml, determined by absorbance using an extinction coefficient ($E_{280}^{1\%}$) of 14.

Buffer:

A buffered stabilizer solution containing 50% (v/v) glycerol.

Storage:

Store between -10 and -20 °C. Product will become viscous but will not freeze. Avoid storage in frost-free freezers. Keep vial tightly capped. Allow product to warm to room temperature and gently mix before use. Avoid exposure to sodium azide as this is an inhibitor of peroxidase activity.

Specificity:

Prior to conjugation, this antibody was specific for Protein C as demonstrated by immunoelectrophoresis and ELISA.

Applications:

Suitable as a source of peroxidase labelled antibodies to PC.

Related Products

GAPC-IG	Goat Anti-Human PC, IgG from antiserum
GAPC-AP	Goat Anti-Human PC, affinity purified IgG
SAPC-IG	Sheep Anti-Human PC, IgG from antiserum
SAPC-AP	Sheep Anti-Human PC, affinity purified IgG
MAPC-IG	Murine Anti-Human PC, IgG from ascites
PC-EIA	Paired antibody ELISA of PC, 5 x 96 wells
APCAT-EIA	ELISA of APC-α ₁ AT complex, 5 x 96 wells
APCPCI-EIA	ELISA of APC-PCI complex, 5 x 96 wells
PC-DP	Human plasma deficient in PC, immune depleted

Visit our site (www.enzymeresearch.co.uk).

Enzyme Research Laboratories Ltd
Glamorgan House . Charter Court . Enterprise Park .
Swansea . SA7 9DD . UK.
Tel: +44 (0)1792 686070
Email: info@enzymeresearch.co.uk
Website: www.enzymeresearch.co.uk